

Fodder crops

Seed treatment		Processing during the growing season		
Preparation of working solution	Working solution consumption	Preparation of working solution	Working solution consumption	Spraying phases
Corn (for silage, green forage)				
150-200 ml REFITON + 10 l of water	10 l for 1 ton of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> . The germination phase - the appearance of the 3rd leaf. . Panicle-flowering phase.
Clover, alfalfa				
150-200ml REFITON + 10 liters of water	10 l for 1 ton of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> . Germination phase. . Plant regrowth phase. . Stalking phase.
Cereals (hedgehog combined, meadow fescue)				
150-200ml REFITON + 10 liters of water	10 l for 1 ton of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> . Germination phase. . Plant regrowth phase. . Stalking phase.
Vico-oat mixture				
150-200ml REFITON + 10 liters of water	10 l for 1 ton of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> . The germination phase is the beginning of plant regrowth. . 15 days after the first. . 15 days after the second.
Natural hayfields and pasture				
		400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> . The germination phase is the beginning of plant regrowth. . 15 days after the first. . 15 days after the second.

* - depending on the type of sprayer