

Potatoes and vegetables and melons

Recommendations for use

Seed treatment		Processing during the growing season		
Preparation of working solution	Working solution consumption	Preparation of working solution	Working solution consumption	Spraying phases
Potato				
150-200 ml REFITON + 40 l of water	40 l for 1 ton of tubers	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> 6-8 leaf phase. The budding phase is the beginning of flowering.
Beetroot				
150-200ml REFITON +15 l of water	15 l for 1 ton of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> Phase 2-3 true leaves. The phase of closing plant leaves in rows.
Tomatoes, cucumbers				
150-200ml REFITON + 10 liters of water	10 l for 10 kg of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> Phase 3-4 of true leaves. With an interval of 12-15 days. With an interval of 12-15 days.
Cabbage				
150-200ml REFITON + 10 liters of water	10 l for 10 kg of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> 2-3 days after planting in the ground. The phase of the leaf whorl is the set of heads. 10-12 days after the second.
Watermelon, melon, pumpkin				
150-200ml REFITON +20 l of water	10 l for 10 kg of seeds	400 ml REFITON for 50-300 * l of water	50-300 * l per 1 ha	<ul style="list-style-type: none"> Lash formation phase. 15-20 days after the first one.

* - depending on the type of sprayer